



CAREER ENDEAVOUR

ACADEMY PRIVATE LIMITED

Best Institute for IIT-JAM|CSIR-UGC-NET/JRF|GATE

IIT-JAM CHEMISTRY-(CY) 2016

TEST : ATOMIC STRUCTURE

Time : 00:50 Hour

Date : 04-07-2015

M.M. : 60

Instructions:

- **Part-A** contains 10 Multiple Choice Questions (MCQ). Each question has 4 choices (a), (b), (c) and (d), for its answer, out of which **ONLY ONE** is correct. For each correct answer you will be awarded **3 marks**. For each incorrect answered **1 mark** will be deducted.
- **Part-B** contains 05 Multiple Select Questions (MSQ). Each question has 4 choices (a), (b), (c) and (d) for its answer, out of which **ONE or MORE than ONE** is/are correct. For each correct answer you will be awarded **2 marks**, there is no negative marking in this section.
- **Part-C** contains 10 Numerical Answer Type (NAT) questions which contain **2 Marks** for each, and there is no negative marking.

PART- A

1. A particle is moving in one dimensional box between $x = a$ and $x = b$. The potential energy is such that the particle can not be outside these limits and the wave function in between is $\psi = \frac{A}{x}$. The average value of x is:

(a) $\sqrt{\frac{ab}{b-a}}$ (b) $\sqrt{\frac{ab}{b-a}} \ln \frac{b}{a}$ (c) $\frac{ab}{b-a} \ln \frac{b}{a}$ (d) $\sqrt{\frac{ab}{b-a}} \ln \frac{a}{b}$
2. An electron is confined in a 1-D box of 1 nm long. How many energy levels are there with energy less than 10 eV is:

(a) 5 (b) 10 (c) 15 (d) 16
3. Identify which of the following operators is not hermitian ?

(a) $\frac{\hbar}{i} \frac{d}{dx}$ (b) $i \frac{d^2}{dx^2}$ (c) $\frac{d^2}{dx^2}$ (d) x^2
4. The angle between the angular momentum $l = 2$ and the z-axis for $m_l = 2$ is:

(a) 35.3 (b) 65.9 (c) 114.1 (d) 144.7
5. A hydrogenic orbital with radial function of the form $r^a \exp[-\beta r]$ and ϕ -part as $\exp[-3i\phi]$ corresponds to

(a) $n > 4, l > 3, m = 3$ (b) $n = 4, l = 3, m = -3$
 (c) $n = 4, l > 3, m = 3$ (d) $n > 4, l = 3, m = -3$
6. A particle is in a one-dimensional box with a potential V_0 inside the box and infinite outside. An energy state corresponding to $n = 0$ (n : quantum number) is not allowed because

(a) the total energy becomes zero
 (b) the average momentum becomes zero
 (c) the wave function becomes zero everywhere
 (d) the potential $V_0 \neq 0$



7. An eigenstate of energy satisfies $H\psi_n = E_n\psi_n$. In the presence of an extra constant potential V_0
- (a) both E_n and ψ_n will change (b) both E_n and average kinetic energy will change
 (c) only E_n will change, but not ψ_n (d) only ψ_n will change, but not E_n .
8. Compare the difference of energies of the first excited and ground states of a particle confined in (i) a 1-d box (Δ_1), (ii) a 2-d square box (Δ_2) and (iii) a 3-d cubic box (Δ_3). Assume the length of each of the boxes is the same. The correct relation between the energy differences Δ_1 , Δ_2 and Δ_3 for the three states is
- (a) $\Delta_1 > \Delta_2 > \Delta_3$ (b) $\Delta_1 = \Delta_2 = \Delta_3$ (c) $\Delta_3 > \Delta_2 > \Delta_1$ (d) $\Delta_3 > \Delta_1 > \Delta_2$
9. The correct statement about both the average value of position ($\langle x \rangle$) and momentum ($\langle p \rangle$) of a 1-d harmonic oscillator wavefunction is
- (a) $\langle x \rangle \neq 0$ and $\langle p \rangle \neq 0$ (b) $\langle x \rangle = 0$ but $\langle p \rangle \neq 0$
 (c) $\langle x \rangle = 0$ and $\langle p \rangle = 0$ (d) $\langle x \rangle \neq 0$ but $\langle p \rangle = 0$
10. The value of the commutator $[x, [x, p_x]]$ is
- (a) $i\hbar x$ (b) $-i\hbar$ (c) $i\hbar$ (d) 0

PART-B

11. Analyse the function given, $\psi_{n,\ell,m} \propto \rho^2 e^{-\rho/3} \sin \theta \cos \theta \cos \phi$
The correct statement is
- (a) Angular node corresponding to this function = 3 and radial node corresponding to this function = 0
 (b) The z-component of angular momentum corresponding to the above function is $\frac{h}{2\pi}$
 (c) Total angular momentum corresponding to this function $\frac{3h^2}{2\pi^2}$ and energy corresponding to this function is -0.055 a.u.
 (d) Average value of $\langle r \rangle$ corresponding to above function is $12a_0$.
12. Consider the statement. The **correct** statements are
- (a) In the Bohr atom the angular momentum always had a non-zero value. While in the modern theory the angular momentum is zero in S state only
 (b) In Schrodinger model n has nothing directly to do with angular momentum, while on Bohr model n was measure first of the angular momentum of the system.
 (c) If a hydrogen atom in the ground 1s state absorb radiation, then it must be finally in 2s state.
 (d) The virial theorem corresponding to H-atom is $2 \times \text{total energy} = - \langle \text{potential energy} \rangle$
13. The **incorrect** statement is
- (a) The degeneracy corresponding to Li^{++} species having energy $-\frac{e^2}{72\pi \epsilon_0 a_0}$ is 9
 (b) The degeneracy corresponding to rotating a particle of mass 'm' having energy $\frac{5h^2}{2\pi^2 mr^2}$ is 7
 (c) The energy corresponding to particle confined to move in 2D square box having energy $\frac{25h^2}{4\ell^2}$ is 2
 (d) The negative value of energy eigenvalue $E \left(= \frac{-13.6z^2}{n^2} \right)$ corresponds to scattering states in which there is not enough energy for the electron to escape from the nucleus



14. $\psi = N r (6 - Z r) e^{-Zr/3} \cos \theta$, is a proposed hydrogenic wavefunction, where Z = atomic number, r = radial distance from the nucleus, θ = azimuthal angle, N is a constant. The **CORRECT** statement about ψ is
- $\psi = 0$ in the xy -plane
 - two radial nodes are present in ψ
 - one angular node is present in ψ
 - the size of the orbital decreases with increase in atomic number
15. Among the following statements, which statements **TRUE** for a Hermitian operator are :
- The eigen values are real
 - the eigen functions are orthogonal
 - The represent observables
 - The scalar product of two different eigen function is equal to one

PART- C

16. The wavelength associated with an electron moving with a velocity of 10^3 m sec^{-1} _____ (-10^{-7} m).
17. The degeneracy of the energy level $\frac{12h^2}{8ml^2}$ of a particle in a three dimensional cube of the length 'a' is _____
18. In the units of $\frac{h^2}{8ml^2}$, the energy difference between levels corresponding to 3 to 2 node eigen functions for a particle of mass m in a one dimensional box of length 'l' is _____
19. For unnormalized wave-function, $\psi(r, \theta, \phi) = \sin \theta \cos \phi \left(\frac{2r}{a_0} - \left(\frac{r}{a_0} \right)^2 \right) \exp \left(-\frac{r}{a_0} \right)$, the number of radial node(s) is _____
20. The wavelength of an electron accelerated through a potential of 100 kV is _____ (\AA)
21. The probability of finding the particle in one dimensional box of length L in the region between $\frac{L}{4}$ and $\frac{3L}{4}$ for quantum number $n = 2$ is _____ %
22. The ground state translational energy of a particle in one-dimensional box of 300pm length is about 4 eV. Suppose that the same particle is moving in a three-dimensional cubic box of 100 pm on the side. The ground state energy of the particle in the three-dimensional box is _____ (eV).
23. A cubic box of edge-length 1.2 nm contains 10 electrons. Applying the simple particle in a box theory, the value of ΔE for the first excited state of this system is _____ (10^{-20} J)
24. For a particle of mass m confined in a box of length L , assume $\Delta x = L$. Assume further that $\Delta p(\text{min}) = \langle p^2 \rangle^{1/2}$. Use the uncertainty principle to obtain an estimate of the energy of the particle. The value will be _____ $\left(\frac{h^2}{mL^2} \right)$.
25. The ground state energy of hydrogen atom is -13.598 eV . The expectation values of kinetic energy, $\langle T \rangle$ is _____ (in units of eV).





CAREER ENDEAVOUR

ACADEMY PRIVATE LIMITED

Best Institute for IIT-JAM, NET/JRF & GATE Exams

IIT-JAM CHEMISTRY-2016

Date : 04-07-2015

TEST : ATOMIC STRUCTURE

ANSWER KEY

Part - A

1. (c) 2. (a) 3. (b) 4. (a) 5. (d) 6. (c) 7. (c)
8. (b) 9. (c) 10. (d)

Part - B

11. (b,c,d) 12. (a,b) 13. (a,b,c,d) 14. (a,c,d) 15. (a,b,c)

Part - C

16. (7.25) 17. (1) 18. (7) 19. (1) 20. (0.037) 21. (50)
22. (108) 23. (8.366) 24. (0.125) 25. (13.598)

