

PAPER : IIT-JAM 2006
MATHEMATICS-MA

Q.1-Q.15: Only one option is correct for each question. Each question carries (+6) marks for correct answer and (-2) marks for incorrect answer.

1. $\lim_{n \rightarrow \infty} \frac{2^{n+1} + 3^{n+1}}{2^n + 3^n}$ equals
(a) 3 (b) 2 (c) 1 (d) 0
 2. Let $f(x) = (x - 2)^{17}(x + 5)^{24}$. Then
(a) f does not have a critical point at 2 (b) f has a minimum at 2
(c) f has a maximum at 2 (d) f has neither a minimum nor a maximum at $x = 2$
 3. Let $f(x, y) = x^5 y^2 \tan^{-1}\left(\frac{y}{x}\right)$. Then $x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y}$ equals
(a) $2f$ (b) $3f$ (c) $5f$ (d) $7f$
 4. Let G be the set of all irrational numbers. The interior and the closure of G are denoted by G^0 and \bar{G} , respectively. Then
(a) $G^0 = \phi, \bar{G} = G$ (b) $G^0 = \mathbb{R}, \bar{G} = \mathbb{R}$ (c) $G^0 = \phi, \bar{G} = \mathbb{R}$ (d) $G^0 = G, \bar{G} = \mathbb{R}$
 5. Let $f(x) = \int_{\sin x}^{\cos x} e^{-t^2} dt$, then $f'(\pi/4)$ equals
(a) $\sqrt{1/e}$ (b) $-\sqrt{2/e}$ (c) $\sqrt{2/e}$ (d) $-\sqrt{1/e}$
 6. Let C be the circle $x^2 + y^2 = 1$ taken in anti-clockwise sense. Then the value of the integral $\int_C [(2xy^3 + y)dx + (3x^2y^2 + 2x)dy]$ equals
(a) 1 (b) $\pi/2$ (c) π (d) 0
 7. Let r be the distance of a point $P(x, y, z)$ from the origin O . The ∇r is a vector
(a) orthogonal to \overline{OP}
(b) normal to the level surface of r at P
(c) normal to the surface of revolution generated by OP about x -axis
(d) normal to the surface of revolution generated by OP about y -axis.
 8. Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by $T(x_1, x_2, x_3) = (x_1 - x_2, x_1 - x_2, 0)$. If $N(T)$ and $R(T)$ denote the null space and range space of T respectively, then
(a) $\dim N(T) = 2$ (b) $\dim R(T) = 2$ (c) $R(T) = N(T)$ (d) $N(T) \subset R(T)$
 9. Let S be a closed surface for which $\iint_S \vec{r} \cdot \hat{n} d\sigma = 1$. Then the volume enclosed by the surface is
(a) 1 (b) $1/3$ (c) $2/3$ (d) 3
-



10. If $(c_1 + c_2 \ln x)/x$ is the general solution of the differential equation

$$x^2 \frac{d^2 y}{dx^2} + kx \frac{dy}{dx} + y = 0, \quad x > 0, \text{ then } k \text{ equals}$$

- (a) 3 (b) -3 (c) 2 (d) -1
11. If A and B are 3×3 real matrices such that $\text{rank}(AB) = 1$, then $\text{rank}(BA)$ **cannot** be
- (a) 0 (b) 1 (c) 2 (d) 3
12. The differential equation representing the family of circles touching y-axis at the origin is
- (a) linear and of first order (b) linear and of second order
(c) nonlinear and of first order (d) nonlinear and of second order
13. Let G be a group of order 7 and $\phi(x) = x^4, x \in G$. Then ϕ is
- (a) not one-one (b) not onto
(c) not a homomorphism (d) one-one, onto and a homomorphism
14. Let R be the ring of all 2×2 matrices with integer entries. Which of the following subsets of R is an integral domain?
- (a) $\left\{ \begin{pmatrix} 0 & x \\ y & 0 \end{pmatrix} : x, y \in \mathbf{Z} \right\}$ (b) $\left\{ \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix} : x, y \in \mathbf{Z} \right\}$
(c) $\left\{ \begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix} : x \in \mathbf{Z} \right\}$ (d) $\left\{ \begin{pmatrix} x & y \\ y & z \end{pmatrix} : x, y, z \in \mathbf{Z} \right\}$
15. Let $f_n(x) = n \sin^{2n+1} x \cos x$. Then the value of $\lim_{n \rightarrow \infty} \int_0^{\pi/2} f_n(x) dx - \int_0^{\pi/2} \left(\lim_{n \rightarrow \infty} f_n(x) \right) dx$ is
- (a) $\frac{1}{2}$ (b) 0 (c) $-\frac{1}{2}$ (d) $-\infty$
16. (a) Test the convergence of the series $\sum_{n=1}^{\infty} \frac{n^n}{n! 3^n}$ (6)
- (b) Show that $\ln(1 + \cos x) \leq \ln 2 - \frac{x^2}{4}$ for $0 \leq x \leq \frac{\pi}{2}$. (9)
17. Find the critical points of the function $f(x, y) = x^3 + y^2 - 12x - 6y + 40$. Test each of these for maximum and minimum. (15)
18. (a) Evaluate $\iint_R x e^{y^2} dx dy$, where R is the region bounded by the lines $x = 0, y = 1$ and the parabola $y = x^2$. (6)
- (b) Find the volume of the solid bounded above by the surface $z = 1 - x^2 - y^2$ and below by the plane $z = 0$. (9)

19. Evaluate the surface integral $\iint_S x(12y - y^4 + z^2)d\sigma$, where the surface S is represented in the form $z = y^2, 0 \leq x \leq 1, 0 \leq y \leq 1$. (15)
20. Using the change of variables evaluate $\iint_R xy \, dx \, dy$, where the region R is bounded by the curves $xy = 1, xy = 3, y = 3x$ and $y = 5x$ in the first quadrant. (15)
21. (a) Let u and v be the eigenvectors of A corresponding to the eigenvalues 1 and 3 respectively. Prove that $u + v$ is not an eigenvector of A . (6)
- (b) Let A and B be real matrices such that the sum of each row of A is 1 and the sum of each row of B is 2. Then show that 2 is an eigenvalue of AB . (9)
22. Suppose W_1 and W_2 are subspace of \mathbb{R}^4 spanned by $\{(1, 2, 3, 4), (2, 1, 1, 2)\}$ and $\{(1, 0, 1, 0), (3, 0, 1, 0)\}$ respectively. Find a basis of $W_1 \cap W_2$. Also find a basis of $W_1 + W_2$ containing $\{(1,0,1,0), (3,0,1,0)\}$. (15)
23. Determine y_0 such that the solution of the differential equation $y' - y = 1 - e^{-x}, y(0) = y_0$ has a finite limit as $x \rightarrow \infty$. (15)
24. Let $\phi(x, y, z) = e^x \sin y$. Evaluate the surface integral $\iint_S \frac{\partial \phi}{\partial n} d\sigma$, where S is the surface of the cube $0 \leq x \leq 1, 0 \leq y \leq 1, 0 \leq z \leq 1$ and $\frac{\partial \phi}{\partial n}$ is the directional derivative of ϕ in the direction of the unit outward normal to S . Verify the divergence theorem. (15)
25. Let $y = f(x)$ be a twice continuously differential function on $(0, \infty)$ satisfying $f(1) = 1$ and $f'(x) = \frac{1}{2} f\left(\frac{1}{x}\right), x > 0$. Form the second order differential equation satisfied by $y = f(x)$, and obtain its solution satisfying the given conditions. (15)
26. Let $G = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{Z} \right\}$ be the group under matrix addition and H be the subgroup of G consisting of matrices with even entries. Find the order of the quotient group G/H . (15)
27. Let $f(x) = \begin{cases} x^2 & 0 \leq x \leq 1 \\ \sqrt{x} & x > 1. \end{cases}$
Show that f is uniformly continuous on $[0, \infty)$. (15)
28. Find $M_n = \max_{x \geq 0} \left\{ \frac{x}{n(1 + nx^3)} \right\}$, and hence prove that the series
$$\sum_{n=1}^{\infty} \frac{x}{n(1 + nx^3)}$$
 is uniformly convergent on $[0, \infty)$. (15)

29. Let R be the ring of polynomials with real coefficients under polynomial addition and polynomial multiplication. Suppose

$$I = \{p \in R : \text{sum of the coefficients of } p \text{ is zero}\}.$$

Prove that I is a maximal ideal of R .

(15)

***** END *****

