

SECTION-A

Multiple Choice Questions (MCQ)

Q.1 – Q.10 carry ONE mark each.

- The tangent plane to the surface $z = \sqrt{x^2 + 3y^2}$ at $(1, 1, 2)$ is given by
 - $x - 3y + z = 0$
 - $x + 3y - 2z = 0$
 - $2x + 4y - 3z = 0$
 - $3x - 7y + 2z = 0$
- Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a scalar field, $\vec{v} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ and let $\vec{a} \in \mathbb{R}^3$ be a constant vector. If \vec{r} represents the position vector $x\hat{i} + y\hat{j} + z\hat{k}$, then which one of the following is FALSE?
 - $\text{curl}(f\vec{v}) = \text{grad}(f) \times \vec{v} + f \text{curl}(\vec{v})$
 - $\text{div}(\text{grad}(f)) = \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) f$
 - $\text{curl}(\vec{a} \times \vec{r}) = 2|\vec{a}|\vec{r}$
 - $\text{div}\left(\frac{\vec{r}}{|\vec{r}|^3}\right) = 0$, for $\vec{r} \neq \vec{0}$
- In \mathbb{R}^3 , the cosine of the acute angle between the surfaces $x^2 + y^2 + z^2 - 9 = 0$ and $z - x^2 - y^2 + 3 = 0$ at the point $(2, 1, 2)$ is
 - $\frac{8}{5\sqrt{21}}$
 - $\frac{10}{5\sqrt{21}}$
 - $\frac{8}{3\sqrt{21}}$
 - $\frac{10}{3\sqrt{21}}$
- Let $s_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots + \frac{1}{n!}$ for $n \in \mathbb{N}$. Then which one of the following is TRUE for the sequence $\{s_n\}_{n=1}^{\infty}$
 - $\{s_n\}_{n=1}^{\infty}$ converges in \mathbb{Q}
 - $\{s_n\}_{n=1}^{\infty}$ is a Cauchy sequence but does not converge in \mathbb{Q}
 - The sub-sequence $\{s_{k^n}\}_{n=1}^{\infty}$ is convergent in \mathbb{R} only when k is even natural number
 - $\{s_n\}_{n=1}^{\infty}$ is not a Cauchy sequence
- Consider the vector space V over \mathbb{R} of polynomial functions of degree less than or equal to 3 defined on \mathbb{R} . Let $T : V \rightarrow V$ be defined by $(Tf)(x) = f(x) - xf'(x)$. Then the rank of T is
 - 1
 - 2
 - 3
 - 4



6. Let a be a positive real number. If f is a continuous and even function defined on the interval $[-a, a]$, then

$\int_{-a}^a \frac{f(x)}{1+e^x} dx$ is equal to

- (a) $\int_0^a f(x) dx$ (b) $2 \int_0^a \frac{f(x)}{1+e^x} dx$ (c) $2 \int_0^a f(x) dx$ (d) $2a \int_0^a \frac{f(x)}{1+e^x} dx$

7. In \mathbb{R}^2 , the family of trajectories orthogonal to the family of asteroids $x^{2/3} + y^{2/3} = a^{2/3}$ is given by

- (a) $x^{4/3} + y^{4/3} = c^{4/3}$ (b) $x^{4/3} - y^{4/3} = c^{4/3}$
 (c) $x^{5/3} - y^{5/3} = c^{5/3}$ (d) $x^{2/3} - y^{2/3} = c^{2/3}$

8. If (v_1, v_2, v_3) is a linearly independent set of vectors in a vector space over \mathbb{R} , then which one of the following sets is also linearly independent?

- (a) $\{v_1 + v_2 - v_3, 2v_1 + v_2 + 3v_3, 5v_1 + 4v_2\}$
 (b) $\{v_1 - v_2, v_2 - v_3, v_3 - v_1\}$
 (c) $\{v_1 + v_2 - v_3, v_2 + v_3 - v_1, v_3 + v_1 - v_2, v_1 + v_2 + v_3\}$
 (d) $\{v_1 + v_2, v_2 + 2v_3, v_3 + 3v_1\}$

9. Which one of the following is TRUE?

- (a) \mathbb{Z}_n is cyclic if and only if n is prime
 (b) Every proper sub-group of \mathbb{Z}_n is cyclic
 (c) Every proper sub-group of S_4 is cyclic
 (d) If every proper sub-group of a group is cyclic, then the group is cyclic

10. Let $a_n = \frac{b_{n+1}}{b_n}$, where $b_1 = 1$, $b_2 = 1$ and $b_{n+2} = b_n + b_{n+1}$, $n \in \mathbb{N}$. Then $\lim_{n \rightarrow \infty} a_n$ is

- (a) $\frac{1-\sqrt{5}}{2}$ (b) $\frac{1-\sqrt{3}}{2}$ (c) $\frac{1+\sqrt{3}}{2}$ (d) $\frac{1+\sqrt{5}}{2}$

Q.11 – Q.30 carry TWO marks each.

11. For $x \in \mathbb{R}$, let $f(x) = \begin{cases} x^3 \sin(\frac{1}{x}), & x \neq 0 \\ 0, & x = 0 \end{cases}$. Then which one of the following is FALSE?

- (a) $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 0$



- (b) $\lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 0$
- (c) $\frac{f(x)}{x^2}$ has infinitely many maxima and minima on the interval $(0, 1)$
- (d) $\frac{f(x)}{x^4}$ is continuous at $x = 0$ but not differentiable at $x = 0$
12. If $\vec{F}(x, y) = (3x - 8y)\hat{i} + (4y - 6xy)\hat{j}$ for $(x, y) \in \mathbb{R}^2$, then $\oint_C \vec{F} \cdot d\vec{r}$, where C is the boundary of the triangular region bounded by lines $x = 0$, $y = 0$ and $x + y = 1$ oriented in the anti-clockwise directions, is
- (a) $\frac{5}{2}$ (b) 3 (c) 4 (d) 5
13. Let $f(x, y) = \begin{cases} \frac{xy}{(x^2 + y^2)^\alpha}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$. Then which one of the following is TRUE for f at the point $(0, 0)$?
- (a) For $\alpha = 1$, f is continuous but not differentiable
- (b) For $\alpha = \frac{1}{2}$, f is continuous and differentiable
- (c) For $\alpha = \frac{1}{4}$, f is continuous and differentiable
- (d) For $\alpha = \frac{3}{4}$, f is neither continuous nor differentiable
14. Let $a, b, c \in \mathbb{R}$. Which of the following values of a, b, c do NOT result in the convergence of the series $\sum_{n=3}^{\infty} \frac{a^n}{n^b (\log_e n)^c}$?
- (a) $|a| < 1, b \in \mathbb{R}, c \in \mathbb{R}$ (b) $a = 1, b > 1, c \in \mathbb{R}$
- (c) $a = 1, b \geq 0, c < 1$ (d) $a = -1, b \geq 0, c > 0$
15. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function and let J be a bounded open interval in \mathbb{R} . Define $W(f, J) = \sup\{f(x) \mid x \in J\} - \inf\{f(x) \mid x \in J\}$. Which one of the following is FALSE?
- (a) $W(f, J_1) \leq W(f, J_2)$ if $J_1 \subset J_2$
- (b) If f is a bounded function in J and $J \supset J_1 \supset J_2 \dots \supset J_n \supset \dots$ such that the length of the interval J_n tends to 0 as $n \rightarrow \infty$, then $\lim_{n \rightarrow \infty} W(f, J_n) = 0$

- (c) If f is discontinuous at a point $a \in J$, then $W(f, J) \neq 0$
- (d) If f is continuous at a point $a \in J$, then for any given $\epsilon > 0$ there exists an interval $I \subset J$ such that $W(f, I) < \epsilon$
16. Let U, V and W be finite dimensional real vector spaces, $T:U \rightarrow V$, $S:V \rightarrow W$ and $P:W \rightarrow U$ be linear transformations. If $\text{range}(ST) = \text{nullspace}(P)$, $\text{nullspace}(ST) = \text{range}(P)$ and $\text{rank}(T) = \text{rank}(S)$, then which one of the following is TRUE?
- (a) nullity of $T = \text{nullity of } S$
- (b) dimension of $U \neq \text{dimension of } W$
- (c) If dimension of $V = 3$, dimension of $U = 4$, then P is not identically zero
- (d) If dimension of $V = 4$, dimension of $U = 3$, and T is one-one, then P is identically zero
17. Let $a_n = \frac{(-1)^n}{\sqrt{1+n}}$ and let $c_n = \sum_{k=0}^n a_{n-k} a_k$, where $n \in \mathbb{N} \cup \{0\}$. Then which one of the following is TRUE?
- (a) Both $\sum_{n=0}^{\infty} a_n$ and $\sum_{n=1}^{\infty} c_n$ are convergent
- (b) $\sum_{n=0}^{\infty} a_n$ is convergent but $\sum_{n=1}^{\infty} c_n$ is not convergent
- (c) $\sum_{n=1}^{\infty} c_n$ is convergent but $\sum_{n=0}^{\infty} a_n$ is not convergent
- (d) Neither $\sum_{n=0}^{\infty} a_n$ nor $\sum_{n=1}^{\infty} c_n$ is convergent
18. Let G be a group satisfying the property that $f:G \rightarrow \mathbb{Z}_{221}$ is a homomorphism implies $f(g) = 0, \forall g \in G$. Then a possible group G is
- (a) \mathbb{Z}_{21} (b) \mathbb{Z}_{51} (c) \mathbb{Z}_{91} (d) \mathbb{Z}_{119}
19. An integrating factor of the differential equation $\left(y + \frac{1}{3}y^3 + \frac{1}{2}x^2\right)dx + \frac{1}{4}(x + xy^2)dy = 0$ is
- (a) x^2 (b) $3\log_e x$ (c) x^3 (d) $2\log_e x$
20. Let H be the quotient group \mathbb{Q}/\mathbb{Z} . Consider the following statements.
- I. Every cyclic subgroup of H is finite
- II. Every finite cyclic group is isomorphic to a subgroup of H
- Which one of the following holds?
- (a) I is TRUE but II is FALSE (b) II is TRUE but I is FALSE
- (c) both I and II are TRUE (d) neither I nor II is TRUE

21. Let $a_n = \begin{cases} 2 + \frac{(-1)^{\frac{n-1}{2}}}{n}, & \text{if } n \text{ is odd} \\ 1 + \frac{1}{2^n}, & \text{if } n \text{ is even} \end{cases}$, $n \in \mathbb{N}$. Then which one of the following is TRUE?
- (a) $\sup\{a_n | n \in \mathbb{N}\} = 3$ and $\inf\{a_n | n \in \mathbb{N}\} = 1$
- (b) $\liminf(a_n) = \limsup(a_n) = \frac{3}{2}$
- (c) $\sup\{a_n | n \in \mathbb{N}\} = 2$ and $\inf\{a_n | n \in \mathbb{N}\} = 1$
- (d) $\liminf(a_n) = 1$ and $\limsup(a_n) = 3$
22. Let $a_n = n + \frac{1}{n}$, $n \in \mathbb{N}$. Then the sum of the series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{a_{n+1}}{n!}$ is
- (a) $e^{-1} - 1$ (b) e^{-1} (c) $1 - e^{-1}$ (d) $1 + e^{-1}$
23. For $x > \frac{-1}{2}$, let $f_1(x) = \frac{2x}{1+2x}$, $f_2(x) = \log_e(1+2x)$ and $f_3(x) = 2x$. Then which one of the following is TRUE?
- (a) $f_3(x) < f_2(x) < f_1(x)$ for $0 < x < \frac{\sqrt{3}}{2}$ (b) $f_1(x) < f_3(x) < f_2(x)$ for $x > 0$
- (c) $f_1(x) + f_2(x) < \frac{f_3(x)}{2}$ for $x > \frac{\sqrt{3}}{2}$ (d) $f_2(x) < f_1(x) < f_3(x)$ for $x > 0$
24. Consider the group $\mathbb{Z}^2 = \{(a, b) | a, b \in \mathbb{Z}\}$ under component wise addition. Then which of the following is a subgroup of \mathbb{Z}^2 ?
- (a) $\{(a, b) \in \mathbb{Z}^2 | ab = 0\}$ (b) $\{(a, b) \in \mathbb{Z}^2 | 3a + 2b = 15\}$
- (c) $\{(a, b) \in \mathbb{Z}^2 | 7 \text{ divides } ab\}$ (d) $\{(a, b) \in \mathbb{Z}^2 | 2 \text{ divides } a \text{ and } 3 \text{ divides } b\}$
25. A particular integral of the differential equation $y'' + 3y' + 2y = e^{e^x}$ is
- (a) $e^{e^x} e^{-x}$ (b) $e^{e^x} e^{-2x}$ (c) $e^{e^x} e^{2x}$ (d) $e^{e^x} e^x$
26. $a, b \in \mathbb{R}$ and let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a thrice differential function. If $z = e^u f(v)$, where $u = ax + by$ and $v = ax - by$, then which one of the following is TRUE?
- (a) $b^2 z_{xx} - a^2 z_{yy} = 4a^2 b^2 e^u f'(v)$ (b) $b^2 z_{xx} - a^2 z_{yy} = -4e^u f'(v)$
- (c) $b z_x + a z_y = ab z$ (d) $b z_x + a z_y = -ab z$

27. Let $y(x)$ be the solution of the differential equation $\frac{dy}{dx} + y = f(x)$, for $x \geq 0$, $y(0) = 0$, where

$$f(x) = \begin{cases} 2, & 0 \leq x < 1 \\ 0, & x \geq 1 \end{cases}. \text{ Then } y(x) =$$

- (a) $2(1 - e^{-x})$ when $0 \leq x < 1$ and $2(e - 1)e^{-x}$ when $x \geq 1$
 (b) $2(1 - e^{-x})$ when $0 \leq x < 1$ and 0 when $x \geq 1$
 (c) $2(1 - e^{-x})$ when $0 \leq x < 1$ and $2(1 - e^{-1})e^{-x}$ when $x \geq 1$
 (d) $2(1 - e^{-x})$ when $0 \leq x < 1$ and $2e^{1-x}$ when $x \geq 1$

28. Consider the region D in the yz plane bounded by the line $y = \frac{1}{2}$ and the curve $y^2 + z^2 = 1$, where $y \geq 0$.

If the region D is revolved about the z -axis in \mathbb{R}^3 , then the volume of the resulting solid is

- (a) $\frac{\pi}{\sqrt{3}}$ (b) $\frac{2\pi}{\sqrt{3}}$ (c) $\frac{\pi\sqrt{3}}{2}$ (d) $\pi\sqrt{3}$

29. Suppose that $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable functions such that f is strictly increasing and g is strictly decreasing. Define $p(x) = f(g(x))$ and $q(x) = g(f(x))$, $\forall x \in \mathbb{R}$. Then for $t > 0$ the sign of

$$\int_0^t p'(x)(q'(x) - 3)dx \text{ is}$$

- (a) positive (b) negative (c) dependent on t (d) dependent on f and g

30. Let I denote the 4×4 identity matrix. If the roots of the characteristic polynomial of a 4×4 matrix M

$$\text{are } \pm\sqrt{\frac{1 \pm \sqrt{5}}{2}}, \text{ then } M^8 =$$

- (a) $I + M^2$ (b) $2I + M^2$ (c) $2I + 3M^2$ (d) $3I + 2M^2$

SECTION-B

Multiple Select Questions (MSQ)

Q.1 – Q.10 carry TWO marks each.

1. The solution(s) of the differential equation $\frac{dy}{dx} = (\sin 2x)y^{1/3}$ satisfying $y(0) = 0$ is(are)

- (a) $y(x) = 0$ (b) $y(x) = -\sqrt{\frac{8}{27}} \sin^3 x$
 (c) $y(x) = \sqrt{\frac{8}{27}} \sin^3 x$ (d) $y(x) = \sqrt{\frac{8}{27}} \cos^3 x$



2. Let P and Q be two non-empty disjoint subsets of \mathbb{R} . Which of the following is (are) FALSE?
- (a) If P and Q are compact, then $P \cup Q$ is also compact
 (b) If P and Q are not connected, then $P \cup Q$ is also not connected
 (c) If $P \cup Q$ and P are closed, then Q is closed
 (d) If $P \cup Q$ and P are open, then Q is open
3. Let $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ denote the group of non-zero complex numbers under multiplication. Suppose $Y_n = \{z \in \mathbb{C} \mid z^n = 1\}, n \in \mathbb{N}$. Which of the following is (are) subgroup(s) of \mathbb{C}^* ?
- (a) $\bigcup_{n=1}^{100} Y_n$ (b) $\bigcup_{n=1}^{\infty} Y_{2^n}$ (c) $\bigcup_{n=100}^{\infty} Y_n$ (d) $\bigcup_{n=1}^{\infty} Y_n$
4. Let $f: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be defined by $f(x) = x + \frac{1}{x^3}$. On which of the following interval (s) is f one-one?
- (a) $(-\infty, -1)$ (b) $(0, 1)$ (c) $(0, 2)$ (d) $(0, \infty)$
5. Let S be a subset of \mathbb{R} such that 2018 is an interior point of S . Which of the following is (are) TRUE?
- (a) S contains an interval
 (b) There is a sequence in S which does not converge to 2018
 (c) There is an element $y \in S, y \neq 2018$ such that y is also an interior point of S
 (d) There is a point $z \in S$, such that $|z - 2018| = 0.002018$
6. $\vec{F}(x, y, z) = (2x + 3yz)\hat{i} + (3xz + 2y)\hat{j} + (3xy + 2z)\hat{k}$ for $(x, y, z) \in \mathbb{R}^3$, then which among the following is (are) TRUE?
- (a) $\nabla \times \vec{F} = \vec{0}$
 (b) $\oint_C \vec{F} \cdot d\vec{r} = 0$ along any simple closed curve C
 (c) There exists a scalar function $\phi: \mathbb{R}^3 \rightarrow \mathbb{R}$ such that $\nabla \cdot \vec{F} = \phi_{xx} + \phi_{yy} + \phi_{zz}$
 (d) $\nabla \cdot \vec{F} = 0$
7. Let $m, n \in \mathbb{N}, m < n, P \in M_{n \times m}(\mathbb{R}), Q \in M_{m \times n}(\mathbb{R})$. Then which of the following is (are) NOT possible?
- (a) $\text{rank}(PQ) = n$
 (b) $\text{rank}(QP) = m$
 (c) $\text{rank}(PQ) = m$
 (d) $\text{rank}(QP) = \left\lceil \frac{m+n}{2} \right\rceil$, the smallest integer larger than or equal to $\frac{m+n}{2}$

8. Suppose $\alpha, \beta, \gamma \in \mathbb{R}$. Consider the following system of linear equations.
 $x + y + z = \alpha, x + \beta y + z = \gamma, x + y + \alpha z = \beta$. If this system has at least one solution, then which of the following statements is (are) TRUE?
- (a) If $\alpha = 1$ then $\gamma = 1$ (b) If $\beta = 1$ then $\gamma = \alpha$
 (c) If $\beta \neq 1$ then $\alpha = 1$ (d) If $\gamma = 1$ then $\alpha = 1$
9. Suppose f, g, h are permutations of the set $\{\alpha, \beta, \gamma, \delta\}$, where
 f interchanges α and β but fixes γ and δ
 g interchanges β and γ but fixes α and δ
 h interchanges γ and δ but fixes α and β
 Which of the following permutations interchange (s) α and δ but fix(es) β and γ ?
 (a) $f \circ g \circ h \circ g \circ f$ (b) $g \circ h \circ f \circ h \circ g$ (c) $g \circ f \circ h \circ f \circ g$ (d) $h \circ g \circ f \circ g \circ h$
10. Which of the following subsets of \mathbb{R} is (are) connected?
 (a) $\{x \in \mathbb{R} \mid x^2 + x > 4\}$ (b) $\{x \in \mathbb{R} \mid x^2 + x < 4\}$
 (c) $\{x \in \mathbb{R} \mid |x| < |x - 4|\}$ (d) $\{x \in \mathbb{R} \mid |x| > |x - 4|\}$

SECTION-C

Numerical Answer Type (NAT)

Q.1 – Q.10 carry ONE mark each.

1. Let $f(x) = \sum_{n=0}^{\infty} (-1)^n x(x-1)^n$ for $0 < x < 2$. Then the value of $f\left(\frac{\pi}{4}\right)$ is _____
2. Let $f(x, y) = \sqrt{x^3 y} \sin\left(\frac{\pi}{2} e^{\left(\frac{y-1}{x}\right)}\right) + xy \cos\left(\frac{\pi}{3} e^{\left(\frac{x-1}{y}\right)}\right)$ for $(x, y) \in \mathbb{R}^2, x > 0, y > 0$.
 Then $f_x(1, 1) + f_y(1, 1) =$ _____
3. Let $\phi(x, y, z) = 3y^2 + 3yz$ for $(x, y, z) \in \mathbb{R}^3$. Then the absolute value of the directional derivative of ϕ in the direction of the line $\frac{x-1}{2} = \frac{y-2}{-1} = \frac{z}{-2}$, at the point $(1, -2, 1)$ is _____
4. Let $f: [0, \infty) \rightarrow [0, \infty)$ be continuous on $[0, \infty)$ and differentiable on $(0, \infty)$. If $f(x) = \int_0^x \sqrt{f(t)} dt$,
 then $f(6) =$ _____



5. Let $a_n = \frac{(1+(-1)^n)}{2^n} + \frac{(1+(-1)^{n-1})}{3^n}$. Then the radius of convergence of the power series $\sum_{n=1}^{\infty} a_n x^n$ about $x=0$ is _____

6. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be given by $f(x, y) = \begin{cases} \frac{x^2 y(x-y)}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$.

Then $\frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) - \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right)$ at the point $(0, 0)$ is _____

7. Let W_1 be the real vector space of all 5×2 matrices such that the sum of the entries in each row is zero. Let W_2 be the real vector space of all 5×2 matrices such that the sum of the entries in each column is zero. Then the dimension of the space $W_1 \cap W_2$ is _____

8. The coefficient of x^4 in the power series expansion of $e^{\sin x}$ about $x=0$ is _____ (correct upto three decimal places).

9. Let A_6 be the group of even permutations of 6 distinct symbols. Then the number of elements of order 6 in A_6 is _____

10. The order of the element $(123)(246)(456)$ in the group S_6 is _____

Q.11 – Q.20 carry TWO marks each.

11. If $y(x) = v(x) \sec x$ is the solution of $y'' - (2 \tan x) y' + 5y = 0$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$, satisfying $y(0) = 0$ and

$y'(0) = \sqrt{6}$, then $v\left(\frac{\pi}{6\sqrt{6}}\right)$ is _____ (correct upto two decimal places).

12. Suppose x, y, z are positive real numbers such that $x + 2y + 3z = 1$. If M is the maximum value of xyz^2 , then the value of $\frac{1}{M}$ is _____

13. If $\alpha = \int_{\pi/6}^{\pi/3} \frac{\sin t + \cos t}{\sqrt{\sin 2t}} dt$, then the value of $\left(2 \sin \frac{\alpha}{2} + 1\right)^2$ is _____

14. Suppose $Q \in M_{3 \times 3}(\mathbb{R})$ is a matrix of rank 2. Let $T : M_{3 \times 3}(\mathbb{R}) \rightarrow M_{3 \times 3}(\mathbb{R})$ be the linear transformation defined by $T(P) = QP$. Then the rank of T is _____

15. Let $a_k = (-1)^{k-1}$, $s_n = a_1 + a_2 + \dots + a_n$ and $\sigma_n = (s_1 + s_2 + \dots + s_n) / n$, where $k, n \in \mathbb{N}$.

Then $\lim_{n \rightarrow \infty} \sigma_n$ is _____ (correct upto one decimal place)



16. The value of the integral $\int_0^1 \int_x^1 y^4 e^{-xy^2} dy dx$ is _____ (correct upto three decimal places)

17. The area of the parametrized surface

$$S = \{((2 + \cos u) \cos v, (2 + \cos u) \sin v, \sin u) \in \mathbb{R}^3 \mid 0 \leq u \leq \frac{\pi}{2}, 0 \leq v \leq \frac{\pi}{2}\} \text{ is _____}$$

(correct upto two decimal places)

18. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be such that f'' is continuous on \mathbb{R} and $f(0) = 1$, $f'(0) = 0$ and $f''(0) = -1$.

Then $\lim_{x \rightarrow \infty} \left(f \left(\sqrt{\frac{2}{x}} \right) \right)^x$ is _____ (correct upto three decimal places)

19. If $x(t)$ is the solution to the differential equation $\frac{dx}{dt} = x^2 t^3 + xt$, for $t > 0$, satisfying $x(0) = 1$, then the value of $x(\sqrt{2})$ is _____ (correct upto two decimal places)

20. If the volume of the solid in \mathbb{R}^3 bounded by the surfaces $x = -1$, $x = 1$, $y = -1$, $y = 1$, $z = 2$, $y^2 + z^2 = 2$ is $\alpha - \pi$, then $\alpha =$ _____

***** END OF QUESTION PAPER *****





CAREER ENDEAVOUR

Best Institute for IIT-JAM, NET & GATE

IIT-JAM MATHEMATICS - 2018

ANSWER KEY

SECTION-A

- | | | | | |
|---------|---------|---------|---------|---------|
| 1. (b) | 2. (c) | 3. (c) | 4. (b) | 5. (c) |
| 6. (a) | 7. (b) | 8. (d) | 9. (b) | 10. (d) |
| 11. (d) | 12. (b) | 13. (c) | 14. (c) | 15. (b) |
| 16. (c) | 17. (b) | 18. (a) | 19. (c) | 20. (c) |
| 21. (a) | 22. (d) | 23. (*) | 24. (d) | 25. (b) |
| 26. (a) | 27. (a) | 28. (c) | 29. (a) | 30. (c) |

SECTION-B

- | | | | |
|--------------|---------------|-----------|-----------|
| 1. (a, b, c) | 2. (b, c, d) | 3. (b) | 4. (b) |
| 5. (a, b, c) | 6. (a, b, c) | 7. (a, d) | 8. (a, b) |
| 9. (a, d) | 10. (b, c, d) | | |

SECTION-C

- | | | | |
|-----------|-------------|--------------|-------------|
| 1. (1) | 2. (3) | 3. (7) | 4. (9) |
| 5. (2) | 6. (1) | 7. (4) | 8. (-0.125) |
| 9. (0) | 10. (4) | 11. (0.5) | 12. (1152) |
| 13. (3) | 14. (6) | 15. (0.5) | 16. (0.239) |
| 17. (6.5) | 18. (0.367) | 19. (-2.718) | 20. (6) |

