



IIT-JAM PHYSICS BHU-2018

1. A rigid body is constrained to move on a plane. Number of degrees of freedom for it will be
(a) 2 (b) 1 (c) 5 (d) 3
2. Number of generalized coordinates required to describe the motion of a solid cylinder rolling without slipping on an inclined plane is
(a) 5 (b) 2 (c) 3 (d) 4
3. The constraints of a rigid body is
(a) conservative and scleronic (b) conservative and rheonomic
(c) holonomic and rheonomic (d) non-holonomic and scleronic
4. Which one of the following represents the equation of motion for the system described by the Hamiltonian $H(q,p)$?
(a) $\dot{q} = \frac{\partial H}{\partial p}, \dot{p} = \frac{\partial H}{\partial q}$ (b) $-\dot{q} = \frac{\partial H}{\partial p}, \dot{p} = \frac{\partial H}{\partial q}$ (c) $\dot{q} = \frac{\partial H}{\partial p}, \dot{p} = -\frac{\partial H}{\partial q}$ (d) $\dot{q} = \frac{\partial H}{\partial p}, -\dot{p} = \frac{\partial H}{\partial q}$
5. A particle of unit mass moves potential $V(x) = x^3 - 3x + 2$. The angular frequency of small oscillation about the minimum of the potential is
(a) $\sqrt{6}$ (b) $\sqrt{3}$ (c) $\frac{1}{\sqrt{6}}$ (d) $\frac{1}{\sqrt{3}}$
6. A system is described by the Lagrangian $L(r, \theta, \dot{r}, \dot{\theta}) = \frac{1}{2}m\dot{r}^2 + \frac{1}{2}mr^2\dot{\theta}^2 + \frac{1}{r}$
Which one of the following is not true?
(a) Total energy of the system is conserved (b) Angular momentum of the system is conserved
(c) θ is cyclic coordinate (d) Linear momentum of system is conserved
7. If q_1 and q_2 are generalized coordinates and p_1 and p_2 are corresponding generalized momenta, then the Poisson bracket $\{q_1^2 + q_2^2, 2p_1 + p_2\}$ is
(a) 0 (b) $(q_1 + 2q_2) 2p_1$ (c) $3(q_1^2 + q_2^2)$ (d) $2(2q_1 + q_2)$




8. Lagrangian for simple harmonic oscillator with frequency ω , mass m in one dimension is given by

(a) $\frac{1}{2}m(\dot{x}^2 - \omega^2 x^2)$ (b) $\frac{1}{2}m(\dot{x}^2 + \omega^2 x^2)$ (c) $\frac{1}{2}m(\ddot{x} + \omega^2 x)$ (d) $\frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2$

9. The probability distribution of a variable x in the range $-\infty$ and $+\infty$ is given by $P(x) = 10e^{-(2x^2 - 4x - 6)}$. The maximum probability will correspond to

(a) $x = 1$ (b) $x = 0$ (c) $x = 3$ (d) $x = -1$

10. The phase space trajectory of a single particle, falling freely from a height will be



11. Number of microstates for a monoatomic ideal gas with N molecules in a volume V and with total energy E is proportional to

(a) E^N (b) $E^{3N/2}$ (c) $E^{N/2}$ (d) E^{3N}

12. If Q the partition function of a system of particles in canonical ensemble, the average energy of the system is given by

(a) $\bar{E} = \frac{\partial Q}{\partial \beta}$ (b) $\bar{E} = -\frac{\partial Q}{\partial \beta}$ (c) $\bar{E} = \frac{\partial}{\partial \beta} \ln Q$ (d) $\bar{E} = -\frac{\partial}{\partial \beta} \ln Q$

13. Consider a system consisting of two particles each of which can be in any one of three quantum states $0, \epsilon, 2\epsilon$. The number of total configurations when the particles are identical bosons

(a) 9 (b) 6 (c) 5 (d) 3

14. Consider a gas of photons in a cubical container of edge length L and volume $V = L^3$. The mean pressure in terms of mean energy E is given by

(a) $\frac{E}{V}$ (b) $\frac{2}{3} \frac{E}{V}$ (c) $\frac{1}{3} \frac{E}{V}$ (d) 0

15. The statistical system in which both energy and number of particles change are best described by

- (a) micro-canonical ensemble theory
 (b) canonical ensemble theory
 (c) grand-canonical ensemble theory
 (d) both canonical as well as grand-canonical ensemble theory

16. Relative root mean square fluctuation of energy in canonical ensemble theory is

(a) $\propto T^{1/2}$ (b) $\propto T$ (c) $\propto T^2$ (d) $\propto T^{3/2}$

17. Given three isobars, namely; ${}_{11}^{25}\text{Na}$, ${}_{12}^{25}\text{Mg}$ and ${}_{13}^{25}\text{Al}$

- (a) ${}_{11}^{25}\text{Na}$ is stable and the other two are beta emitters
 (b) ${}_{12}^{25}\text{Mg}$ is stable and the other two are beta emitters
 (c) ${}_{13}^{25}\text{Al}$ is stable and the other two are beta emitters
 (d) All nuclei are stable
18. Radiocarbon dating is done by estimating in the specimen
 (a) the ratio of amount of ${}^{14}\text{C}$ to ${}^{12}\text{C}$ still present
 (b) the ratio of amount of ${}^{13}\text{C}$ to ${}^{12}\text{C}$ still present
 (c) the amount of radiocarbon still present
 (d) the amount of ${}^{13}\text{C}$ still present
19. The rate of electron emission from 4 mg of ${}_{80}^{210}\text{Pb}$ with half-life 5 days is
 (a) 1.84×10^{16} (b) 1.84×10^{13} (c) 9.2×10^{11} (d) 9.2×10^{16}
20. A proton with 16 MeV energy is bombarded on ${}_{84}^{216}\text{Po}$ nucleus. The proton is
 (a) scattered (b) reflected back
 (c) captured (d) transmitted through the nucleus.
21. The fission rate of ${}^{235}\text{U}$ to produce energy of 200 MW is
 (a) 6.25×10^{15} fission/sec (b) 6.25×10^{16} fission/sec
 (c) 6.25×10^{18} fission/sec (d) 3.12×10^{20} fission/sec
22. The minimum temperature required to initiate fusion of deuteron and triton is of the order of
 (a) 10^9 K (b) 10^6 K (c) 10^{13} K (d) 10^{15} K
23. The average velocity of nucleons inside the nucleus is of the order of
 (a) 3×10^8 m/s (b) 6×10^7 m/s (c) 3×10^6 m/s (d) 6×10^6 m/s
24. The magnetic dipole and electric quadrupole moment data of deuteron imply that the nuclear force is
 (a) purely central (b) central and spin dependent
 (c) mixture of central and non-central components (d) velocity dependent
25. In a crystal, a lattice plane cuts intercepts of $2a$, $3b$ and $6c$ along the axes, where a , b , c are primitive vectors of the unit cell. The Miller indices of the given plane are
 (a) (321) (b) (231) (c) (123) (d) (213)
26. The total number of Bravais lattices are
 (a) 7 (b) 14 (c) 21 (d) 26
27. Origin of characteristic X-rays is
 (a) photoelectric effect (b) inverse photoelectric effect
 (c) electronic transitions within atoms (d) Compton effect
28. The K_{α} line from Molybdenum has a wavelength of 0.7078 \AA . The wavelength of the K_{α} line of copper

(given atomic number of Molybdenum = 42, atomic number of copper = 29)

- (a) 1.517\AA (b) 1.157\AA (c) 1.175\AA (d) 1.715\AA

29. The relation of the reciprocal basis vector \vec{A} to direct basis vector \vec{a} is given by

- (a) $\vec{A} \cdot \vec{a} = 0$ (b) $\vec{A} \cdot \vec{a} = 2\pi$ (c) $\vec{A} \cdot \vec{a} = \pi$ (d) $\vec{A} \cdot \vec{a} = \frac{\pi}{2}$

30. If current carriers are electrons, the Hall coefficient R_H is

- (a) $R_H = -\frac{1}{ne}$ (b) $R_H = \frac{1}{ne}$ (c) $R_H = \frac{n}{e}$ (d) $R_H = ne$

31. The electron velocity, v_F , at the Fermi surface is

- (a) $\hbar \left(\frac{3\pi^2 N}{V} \right)^{1/3}$ (b) $\frac{\hbar}{m} \left(\frac{3\pi^2 N}{V} \right)^{1/3}$ (c) $\frac{\hbar}{m} \left(\frac{3\pi N}{V} \right)^{1/3}$ (d) $\frac{\hbar}{m} \left(\frac{\pi^2 N}{V} \right)^{1/3}$

32. The Langevin function, $L(\alpha)$ is represented by

- (a) $L(\alpha) = \cot h\alpha$ (b) $L(\alpha) = \left[\cot h\alpha + \frac{1}{\alpha} \right]$
 (c) $L(\alpha) = \left[\cot h\alpha - \frac{1}{\alpha} \right]$ (d) $L(\alpha) = (\cot h\alpha - \alpha)$

33. The curl of the electromagnetic intensity is

- (a) conservative (b) rotational (c) divergent (d) static

34. The direction of propagation of electromagnetic wave is given by

- (a) $\vec{E} \cdot \vec{B}$ (b) \vec{E} (c) $\vec{E} \times \vec{B}$ (d) \vec{B}

35. For good conductors, the skin depth varies inversely with

- (a) ω (b) ω^2 (c) $\sqrt{\omega}$ (d) ω^4

36. The divergence of the curl of a vector field is

- (a) a scalar (b) a vector (c) zero (d) infinity

37. The charge build up in the capacitor is due to which quantity?

- (a) Conduction current (b) Displacement current
 (c) Convection current (d) Direct current

38. In conductors, which condition will be true ?

- (a) $\sigma\omega\varepsilon > 1$ (b) $\frac{\sigma}{(\omega\varepsilon)} > 1$ (c) $\frac{\sigma}{(\omega\varepsilon)} < 1$ (d) $\sigma\omega\varepsilon < 1$

39. The relation between the speed of light permeability and permittivity

- (a) $c = \mu\varepsilon$ (b) $c = \frac{\mu}{\varepsilon}$ (c) $c = \frac{1}{\sqrt{\mu\varepsilon}}$ (d) $c = \frac{1}{\mu\varepsilon}$



40. The phenomenon employed in the waveguide operation is
 (a) reflection (b) refraction (c) total internal reflection (d) absorption
41. The metric of spherical polar coordinates are
 (a) $h_{11} = r, h_{22} = 1, h_{33} = r \sin \theta$ (b) $h_{11} = 1, h_{22} = r, h_{33} = r \sin \theta$
 (c) $h_{11} = r, h_{22} = r, \sin \theta, h_{33} = 1$ (d) $h_{11} = r^2, h_{22} = r^2 \sin^2 \theta, h_{33} = r^2 \sin^2 \theta$
42. Given the transformation $u = x + y, v = x - y$ and $du dv = k dx dy$, the value of k is
 (a) 1 (b) -1 (c) 2 (d) $\frac{1}{2}$
43. Given $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, then $(aI + bA)^n$ is (where I is 2×2 unit vector)
 (a) $a^n I + b^n A$ (b) $a^n I + nab^{n-1} A$ (c) $a^n I + naba$ (d) $a^n I + na^{n-1} bA$
44. Eigenvectors of the matrix $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$ are
 (a) $\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ (b) $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}$ (c) $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ (d) $\frac{1}{\sqrt{2}} \begin{pmatrix} i \\ 1 \end{pmatrix}, \frac{1}{\sqrt{2}} \begin{pmatrix} -i \\ 1 \end{pmatrix}$
45. Given the matrix $\begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$ with one of the eigenvalues equal to -3 , the other two eigenvalues are
 (a) 0, 1 (b) 0, -1 (c) 0, 2 (d) $-3, 5$
46. In the equation $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (q^2 x^2 - m^2) y = 0$
 (a) $x = 0$ and $x = \infty$ are ordinary points
 (b) $x = 0$ and $x = \infty$ are regular singular points
 (c) $x = 0$ is a regular singular point and $x = \infty$ is an irregular singular point
 (d) $x = 0$ and $x = \infty$ are irregular singular points
47. One of the solutions of the equation $(1 - x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + 12y = 0$ is
 (a) $H_4(x)$ (b) $P_3(x)$ (c) $L_4(x)$ (d) $J_4(x)$
48. The Delta function $\delta(x^2 - a^2)$ is equal to
 (a) $\delta(x+a)\delta(x-a)$ (b) $\delta(x+a) + \delta(x-a)$

(c) $\frac{1}{2|a|}[\delta(x+a)+\delta(x-a)]$ (d) $\delta(x+a)-\delta(x-a)$

49. The Fourier coefficients of the function

$$f(x) = \begin{cases} 0 & -L \leq x \leq 0 \\ 1 & 0 \leq x \leq L \end{cases}$$

expanded in Fourier series are

(a) $a_0 = 1, a_n = 0, b_n = \frac{1}{n\pi}[1 - (-1)^n]$ (b) $a_0 = 1, a_n = [1 - (-1)^n], b_n = 0$
 (c) $a_0 = -1, a_n = 0, b_n = 0$ (d) $a_0 = 1, a_n = 1, b_n = \frac{1}{n\pi}[1 - (-1)^n]$

50. The operator $i\hbar \frac{d}{dx} - \hat{x}$ in momentum basis is

(a) $i\hbar \frac{d}{dx} - \hat{p}$ (b) $-i\hbar \frac{d}{d\hat{p}} - \hat{p}$ (c) $-i\hbar \frac{d}{d\hat{p}} + \hat{p}$ (d) $i\hbar \frac{d}{d\hat{p}} + \hat{p}$

51. If a^+ and a are creation and annihilation operators for SHO, then which of the following is not a Hermitian operators

(a) $aa^+ + a^+a$ (b) $aa^+ - a^+a$ (c) $i(a^+ - a)$ (d) $i(a^+ + a)$

52. If the expectation value of the momentum operator in the normalized state $\psi(x)$ is $\langle p \rangle$, then expectation

value of the momentum operator in the state $\psi_1(x) = e^{\frac{i}{\hbar}p_0x} \psi(x)$ will be

(a) $\langle p \rangle + p_0$ (b) $\langle p \rangle - p_0$ (c) $\langle p \rangle$ (d) 0

53. The ground state wave function for a 1-d system described by the potential

$$V(x) = 0 \text{ for } -\frac{L}{2} \leq x \leq \frac{L}{2}$$

$$= \infty \text{ elsewhere}$$

is

(a) $A \cos \frac{\pi x}{L}$ (b) $A \sin \frac{\pi x}{2L}$ (c) $A \sin \frac{\pi x}{L}$ (d) $A \cos \frac{\pi x}{2L}$

54. A simple harmonic oscillator in one dimension has an eigenfunction (of the Hamiltonian) which vanishes 3 times in the interval $0 < x < \infty$ and is odd under parity. The energy eigenvalue for this state is

(a) $\frac{7}{2} \hbar \omega$ (b) $\frac{9}{2} \hbar \omega$ (c) $\frac{13}{2} \hbar \omega$ (d) $\frac{15}{2} \hbar \omega$



55. The raising and lowering of angular momentum operators are defined as $L_{\pm} = L_x \pm iL_y$. The commutator $[L_-, L_z]$ equal to
- (a) $-2\hbar L_-$ (b) $\hbar L_-$ (c) $\hbar L_+$ (d) $-\hbar L_-$
56. The bound state energy for the state $\psi_{5,4,2}(r, \theta, \phi)$ in a H-atom problem is given by
- (a) $-\frac{13.6}{5} \text{ eV}$ (b) $-\frac{13.6}{25} \text{ eV}$ (c) $-13.6 \times 5 \text{ eV}$ (d) $-13.6 \times 25 \text{ eV}$
57. In a H-atom problem if $L_z \psi_{3,2,-2}(r, \theta, \phi) = a\hbar \psi_{3,2,-2}(r, \theta, \phi)$, then a is
- (a) 2 (b) -2 (c) $2\sqrt{3}$ (d) $\sqrt{6}$
58. ψ_1 and ψ_2 are the wave functions of two orthogonal states of a system belonging to the energy eigenvalues E and $-E$, respectively. In a measurement of energy of another state ψ of the system, the expectation value of energy is found to be $\frac{E}{2}$. ψ in terms of ψ_1 and ψ_2
- (a) $\frac{\sqrt{3}}{2} \psi_1 + \frac{1}{2} \psi_2$ (b) $\frac{1}{2}(\psi_1 - \psi_2)$ (c) $\frac{1}{\sqrt{2}}(\psi_1 - \psi_2)$ (d) $\frac{3}{4} \psi_1 + \frac{1}{4} \psi_2$
59. In any Bohr orbit of hydrogen atom, the ratio of the kinetic energy to the potential energy of the electron is
- (a) 1/2 (b) 2 (c) -1/2 (d) -2
60. Considering the nuclear mass finite, the Rydberg constant is maximum for
- (a) hydrogen atom (b) deuterium atom
(c) singly ionized helium atom (d) doubly ionized lithium atom
61. In Sommerfeld's model of atom, the orbits characterized by a particular principle quantum number, n and different azimuthal quantum number $n_0 = 1, 2, 3, \dots, n$ have
- (a) same energy (b) energy in increasing order with n_0
(c) energy in decreasing order with n_0 (d) no associated energy
62. Stern-Gerlach experiment is important because it gives experimental verification of
- (a) Quantization of energy of atom (b) orbital motion of electron
(c) electron spin (d) Sommerfeld model of atom
63. The ratio of orbital magnetic dipole moment μ_l to the orbital angular momentum L of an electron in an orbit is given by
- (a) $\frac{\mu_L}{L} = \frac{\mu_b}{\hbar}$ (b) $\frac{\mu_L}{L} = -\frac{\mu_b}{\hbar}$ (c) $\frac{\mu_L}{L} = -\frac{\mu_b}{2\hbar}$ (d) $\frac{\mu_L}{L} = \frac{\mu_b}{2\hbar}$
64. Larmor frequency is the frequency of precession of

- (a) orbital angular momentum, L about the external magnetic field, B
 (b) spin angular momentum, S about the external magnetic field, B
 (c) total angular momentum, J about the external magnetic field, B
 (d) orbital angular momentum, L about the total angular momentum, J
65. On application of weak magnetic field the sodium line arising due to the transition ${}^2P_{3/2} \rightarrow {}^2S_{1/2}$ will split ideally into
 (a) 2 components (b) 4 components (c) 6 components (d) 10 components
66. The half-width of gain profile of a He-laser is 2×10^{-3} nm. If the length of the cavity is 30 cm, how many longitudinal modes can be excited? The emission wavelength is 6328 \AA
 (a) 1 (b) 2 (c) 3 (d) 4
67. At what temperature; pressure remaining unchanged, will the molecular velocity (root mean square velocity) of hydrogen will be double of its value at NTP?
 (a) $1092 \text{ }^\circ\text{C}$ (b) $819 \text{ }^\circ\text{C}$ (c) $1092 \text{ }^\circ\text{F}$ (d) $819 \text{ }^\circ\text{K}$
68. The mean square speed for the following group of particles (N_i represents the number of particles with speed v_i) will be
- | N_i | v_i (m/sec) |
|-------|---------------|
| 2 | 1.0 |
| 4 | 2.0 |
| 8 | 3.0 |
- (a) 11.33 m/sec (b) $16.43 \text{ m}^2/\text{sec}^2$ (c) 2.67 m/sec (d) 3.36 m/sec
69. The ratio between most probable speed and root mean square speed of a gas molecule is
 (a) $\sqrt{\frac{3}{2}}$ (b) $\sqrt{\frac{3}{8\pi}}$ (c) $\sqrt{\frac{2}{3}}$ (d) $\sqrt{\frac{8}{3\pi}}$
70. The mean free path of molecules of a gas at pressure p and temperature T is 2×10^{-5} cm. The mean free path at pressure $\frac{p}{2}$ and temperature $2T$ will be
 (a) 10^{-5} cm (b) 8×10^{-5} cm (c) 10^{-5} m (d) 8×10^{-5} m
71. For the adiabatic expansion of a blackbody radiation enclosure, which of the following is correct ?
 (a) $V^{1/3}T = \text{constant}$ (b) $V.T = \text{constant}$ (c) $V^{4/3}T = \text{constant}$ (d) $\frac{V}{T} = \text{constant}$
 where V is the volume and T is the temperature of the enclosure.
72. In throttling process
 (a) the enthalpy remains constant (b) temperature remain constant
 (c) Gibbs' free energy remain constant (d) entropy remains constant

73. Which one of the following is correct ?

- (a) $\frac{E_\lambda}{T^4} = \text{constant}$ (b) $\frac{E_\lambda}{T^5} = \text{constant}$ (c) $\frac{E_\lambda}{T^2} = \text{constant}$ (d) $\frac{E_\lambda}{T} = \text{constant}$

where E_λ is spectral emissive power

74. The numerical value of the slope of an isenthalpic curve at any point on a TP- diagram is called

- (a) Joule coefficient (b) Joule-Kelvin coefficient
(c) Van der Waals' constant (d) Virial coefficient

75. Which of the following can be used to produce lowest temperature ?

- (a) Liquefaction of N_2 (b) Liquid He
(c) Adiabatic demagnetization of paramagnetic salts
(d) None of these

76. A mass m of water at T_1K is isobarically and adiabatically mixed with an equal mass of water at T_2K , the entropy change of the universe is

- (a) $2mC_p \ln \frac{(T_1+T_2)/2}{\sqrt{T_1T_2}}$ (b) $2m \ln \frac{(T_1+T_2)/2}{\sqrt{T_1T_2}}$
(c) $2C_p \ln \frac{(T_1+T_2)/2}{\sqrt{T_1T_2}}$ (d) $2mC_p$

Where C_p is specific heat at constant pressure.

77. Thermodynamic equation

$$TdS = C_V dT + \frac{\beta T}{k} dV \text{ is called}$$

- (a) 2nd TdS equation (b) 1st TdS equation (c) 3rd TdS equation (d) None of these

Where terms have their usual meanings

78. Which of the following is correct ?

- (a) $C_P = \left(\frac{\partial V}{\partial T}\right)_P \left(\frac{\partial P}{\partial T}\right)_S$ (b) $C_P = T \left(\frac{\partial V}{\partial T}\right)_P \left(\frac{\partial P}{\partial T}\right)_S$
(c) $C_P = \left(\frac{\partial T}{\partial V}\right)_P \left(\frac{\partial P}{\partial T}\right)_S$ (d) $C_P = T \left(\frac{\partial T}{\partial V}\right)_P \left(\frac{\partial P}{\partial T}\right)_S$

79. In one- dimensional elastic collision of two particles, the ratio of velocities of separation and approach is equal to :

- (a) Coefficient of restitution (b) negative of coefficient of restitution
(c) zero, if collision is perfectly elastic (d) infinite



80. If in an elastic collision, a massive particle against a lighter one at rest :
- it can never bounce back along its original path
 - it may bounce back along its original path
 - the two particles move at right angles to each other after collision
 - None of the above
81. In which of the following conditions, the total linear momentum of the system remains constant ?
- If the resultant external force acting on the system of particles is zero
 - If the resultant external force acting on the system particles is positive
 - If the resultant external force acting on the system particles is $-ve$
 - None of these
82. From the nozzle of rocket, 100 kg of gases are exhausted per sec with a velocity of 1000 m/sec. What force (thrust) does the gas exert on the rocket ?
- 100 kg/sec
 - 10^5 Newton
 - 10^3 Newton
 - 100 Newton
83. If a particle is at rest relative to an observer at rest at the centre of a rotating frame of reference
- centrifugal and Coriolis forces both act
 - only centrifugal force acts
 - only Coriolis force acts
 - None of these
84. The length of a rod, of length 5 m in a frame of reference which is moving with $0.6c$ velocity in a direction making 30° angle with the rod is nearly
- 4.3 m
 - 3.4 m
 - 2.43 m
 - 2.34 m
85. π^+ meson decays into a μ^+ meson and a neutrino with mean lifetime of about 2.5×10^{-8} sec in a frame in which it is at rest. If the velocity of the π^+ meson in the laboratory frame be $0.9c$, then the expected lifetime in this frame is
- 5.7×10^{-8} sec
 - 2.5×10^{-8} sec
 - 3.1×10^{-8} sec
 - None of these
86. The speed of an electron having kinetic energy 2 MeV will be
- 2.93×10^8 m/sec
 - 3×10^8 m/sec
 - 10^8 m/sec
 - 1.5×10^8 m/sec
87. Which of the following relations is correct for modulus of rigidity η , bulk modulus K and Poisson's ratio σ ?
- $\sigma = \frac{K - 2\eta}{6K + 2\eta}$
 - $\sigma = \frac{3K - 2\eta}{K + 2\eta}$
 - $\sigma = \frac{3K - 2\eta}{6K + 2\eta}$
 - $\sigma = \frac{K - 2\eta}{K + 2\eta}$
88. Two wires A and B of the same material and equal lengths but of radii r and $2r$ are soldered coaxially. The free end of B is twisted by an angle Φ . The ratio of the twist at the junction and angle Φ is
- $\frac{16}{1}$
 - $\frac{17}{16}$
 - $\frac{16}{17}$
 - $\frac{1}{16}$
89. Which of the following is true about liquid flow through a capillary tube ?
- The velocity of the liquid layer in contact with the capillary tube is least

- (b) The velocity of the liquid layer in contact with the capillary tube is maximum
 (c) The velocity of the liquid layer at the centre of the capillary tube is minimum
 (d) None of these
90. The depletion region is created by
 (a) ionization (b) diffusion (c) recombination (d) (1), (2) and (3)
91. A silicon diode is in series with a $1\text{ k}\Omega$ resistor and a 5 V battery. If the anode is connected to the +ve battery terminal, the cathode voltage with respect to the negative battery terminal is
 (a) 0.7 V (b) 0.3 V (c) 5.7 V (d) 4.3 V
92. Where will be the position of the Fermi level of the n -type material when $N_D = N_A$?
 (a) E_C (b) E_V (c) $\frac{E_C + E_V}{2}$ (d) None of the above
- where terms have their usual meanings.
93. The mobility of electrons in a material is expressed in unit of
 (a) V/s (b) $\left(\frac{\text{m}^2}{\text{V sec}}\right)$ (c) m^2/s (d) J/K
94. A silicon sample is uniformly doped with 10^{16} phosphorus atoms/ cm^3 and 2×10^{16} boron atoms/ cm^3 . If all the dopants are fully ionized, the material is
 (a) n -type with carrier concentration of $3 \times 10^{16}/\text{cm}^3$
 (b) p -type with carrier concentration of $10^{16}/\text{cm}^3$
 (c) p -type with carrier concentration of $4 \times 10^{16}/\text{cm}^3$
 (d) intrinsic
95. The bias condition for a transistor to be used as a linear amplifier is called
 (a) forward-reverse (b) forward-forward (c) reverse-reverse (d) collector bias
96. Wien-bridge oscillators are based on
 (a) positive feedback (b) negative feedback
 (c) the piezoelectric effect (d) high gain
97. Which of the following is a universal gate ?
 (a) OR gate (b) NOR gate (c) AND gate (d) NOT gate
98. For an ideal dielectric, polarization \vec{P} is given by
 (a) $\vec{P} = \epsilon_0 \vec{E}$ (b) $\vec{P} = (K - 1)\epsilon_0 \vec{E}$ (c) $\vec{P} = (K + 1)\epsilon_0 \vec{E}$ (d) $\vec{P} = \frac{\epsilon_0}{K - 1} \vec{E}$
99. Clausius-Mossotti relation is represented by the equation
 (a) $\alpha = \frac{3\epsilon_0}{n} \frac{K - 1}{K + 2}$ (b) $\alpha = \frac{\epsilon_0}{n} \frac{K - 1}{K + 2}$ (c) $\alpha = \frac{\epsilon_0}{3n} \frac{K - 1}{K + 2}$ (d) $\alpha = \frac{3\epsilon_0}{n} \frac{K + 2}{K - 1}$

where symbols have their usual meanings.



- 100.** The dipole moment of water molecule is 6.2×10^{-30} C-m at 20°C . The polarizability α is
 (a) 3.17×10^{-39} C-m²/V (b) 3.17×10^{-37} C-m²/V
 (c) 3.17×10^{-35} C-m²/V (d) 3.17×10^{-33} C-m²/V
- 101.** Three magnetic vectors are related as
 (a) $\vec{B} = \mu_0 (\vec{M} - \vec{H})$ (b) $\vec{B} = \mu_0 (\vec{M} + \vec{H})$
 (c) $\vec{B} = \mu_0^{-1} (\vec{M} + \vec{H})$ (d) $\vec{B} = \mu_0^{-1} (\vec{M} - \vec{H})$
- 102.** For higher values of temperature, the susceptibility of paramagnetic substances is proportional to
 (a) T (b) $\frac{1}{T}$ (c) T^2 (d) $\frac{1}{T^2}$
- 103.** The loss of energy per hour in the iron core of a transformer, the hysteresis loop of which is equivalent in area to 2500 ergs/cm³ is (given frequency = 50 Hz, density of iron = 7.5 g/cm³, weight of the iron core = 10 kg)
 (a) 5.985×10^2 J (b) 5.985×10^3 J (c) 5.985×10^4 J (d) 5.985×10^5 J
- 104.** A current i is flowing in a toroidal coil of circular cross-section of radius R with N number of turns distributed uniformly over its circumference. If A is the cross-sectional area of the toroid, its self-inductance will be
 (a) $L = \frac{\mu_0 N^2 A}{2\pi R}$ (b) $L = \frac{\mu_0 N^2 A}{\pi R}$ (c) $L = \frac{\mu_0 N^2 A}{4\pi R}$ (d) $L = \frac{\mu_0 N^2 A}{2R}$
- 105.** Two inductors L_1 and L_2 are connected in series. The total inductance L will be
 (a) $L = L_1 + L_2$ (b) $L = L_1 + L_2 + 2M$ (c) $L = L_1 + L_2 + M$ (d) $L = L_1 + L_2 - M$
 where M is mutual inductance of two coils.
- 106.** A circuit containing resistor R_1 , inductor L_1 and capacitor C_1 connected in series gives resonance at the same frequency f as the second similar combination R_2 , L_2 and C_2 . If the two circuits are connected in series, the whole circuit will resonate at the frequency
 (a) $2f$ (b) $\frac{f}{2}$ (c) f (d) $\frac{f}{4}$
- 107.** A capacitor of 250 pF is connected in parallel with a coil having inductance of 16 mH and effective resistance 20Ω . The circuit impedance at resonance is
 (a) $3.2 \times 10^4 \Omega$ (b) $3.2 \times 10^3 \Omega$ (c) $3.2 \times 10^2 \Omega$ (d) $3.2 \times 10^5 \Omega$
- 108.** For dispersive medium, group velocity (v_g) and phase velocity (v_p) are related as
 (a) $v_g = v_p + \lambda \frac{dv_p}{d\lambda}$ (b) $v_g = v_p - \lambda \frac{dv_p}{d\lambda}$ (c) $v_g = v_p + \frac{1}{\lambda} \frac{dv_p}{d\lambda}$ (d) $v_g = v_p - \frac{1}{\lambda} \frac{dv_p}{d\lambda}$
- 109.** Photon of energy 1.02 MeV undergoes Compton scattering through 180° . The energy of the scattered photon is
 (a) 1.02 MeV (b) 0.204 MeV (c) 0.402 MeV (d) 0.240 MeV

- 110.** In Newton's ring experiment, the diameters of the bright rings are proportional to the
- (a) natural number (b) square root of natural numbers
(c) square root of odd numbers (d) odd numbers
- 111.** Thin sheet of a transparent material of refractive index, $\mu = 1.50$ is placed in the path of one of the interfering beams in a biprism experiment with monochromatic source of wavelength, $\lambda = 5000 \text{ \AA}$. The central fringe shifts to a position originally occupied by 10th bright fringe. The thickness of the sheet is
- (a) $1 \times 10^{-5} \text{ m}$ (b) $1.5 \times 10^{-5} \text{ m}$ (c) $2 \times 10^{-5} \text{ m}$ (d) $2.5 \times 10^{-5} \text{ m}$
- 112.** Interference pattern is produced by two point sources S_1 and S_2 on a plane perpendicular to the line joining S_1 and S_2 . What will be the shape of interference fringes ?
- (a) Straight line (b) Circular (c) Parabolic (d) Hyperbolic
- 113.** In order to make a glass plate of refractive index, μ_g non-reflecting over a wide wavelength range around λ , a thin film is deposited on it. The refractive index μ_f and the thickness d of the film should be
- (a) $\mu_f = \sqrt{\mu_g \mu_a}, d = \frac{3\lambda}{4\mu_f}$ (b) $\mu_f = \sqrt{\mu_g \mu_a}, d = \frac{\lambda}{4\mu_f}$
(c) $\mu_f = \sqrt{\mu_g / \mu_a}, d = \frac{\lambda}{4\mu_f}$ (d) $\mu_f = \sqrt{\mu_g / \mu_a}, d = \frac{3\lambda}{4\mu_f}$

Where μ_a is the refractive index of air.

- 114.** When the distance between two mirrors in Michelson interferometer is decreased
- (a) The fringe pattern appears to collapse at the centre
(b) the fringe pattern expands
(c) the fringe pattern remains stable
(d) the shape of the fringe changes
- 115.** The spread of the central maximum in the Fraunhofer diffraction by a single slit is approximately given by.
- (a) $\frac{\lambda}{b} \leq \theta \leq \frac{\lambda}{b}$ (b) $\frac{2\lambda}{b} \leq \theta \leq \frac{2\lambda}{b}$ (c) $\frac{\lambda}{2b} \leq \theta \leq \frac{\lambda}{2b}$ (d) $\frac{\lambda}{b} \leq \theta \leq \frac{\lambda}{2b}$

Where θ is diffraction angle, b is width of the slit and λ is the wavelength of the light used.

- 116.** A 2 mW laser beams of wavelength $\lambda = 6 \times 10^{-5} \text{ cm}$ is focussed on the retina by a human eye lens of focal length $f = 2.5 \text{ cm}$ and pupil diameter 2 mm. The intensity on the retina will be of the order of
- (a) 10^4 W/m^2 (b) 10^6 W/m^2 (c) 10^8 W/m^2 (d) 10^2 W/m^2
- 117.** To increase the resolving power of a grating total number of lines on the grating is increased such that the grating element becomes 2.5λ . How many orders will be seen on the screen?
- (a) First order only
(b) First and second orders only
(c) First, second and third orders only
(d) First, second, third and fourth orders only

118. The radii of the circles of a zone plate is given by $r_n = 0.1\sqrt{n}$ cm. The most intense focal point for wavelength $\lambda = 5 \times 10^{-5}$ cm will be at a distance
- (a) 50 cm (b) 100 cm (c) 150 cm (d) 200 cm
119. What is the minimum thickness of the base of a prism that can just resolve the two lines of sodium light centred at 5890 \AA and 5896 \AA . The given value of refractive index of prism material is 1.6545 at wavelength 6563 \AA and 1.6635 at wavelength 5270 \AA ?
- (a) 8 mm (b) 10 mm (c) 12 mm (d) 14 mm
120. An unpolarized light is incident on a glass plate placed in air at polarizing angle. The reflected light is
- (a) Plane polarized with electric vector perpendicular to the plane of incidence
(b) plane polarized with electric vector parallel to the plane of incidence
(c) partially polarized having more electric field vectors perpendicular to the plane of incidence
(d) partially polarized having more electric field vectors parallel to the plane of incidence.

